Drug Facts

COMPOSITION : Each film coated tablet contains DAPAGLIFLOZIN TABLETS 10 MG	**PHARMACOLOGICAL CLASS :** sodium-glucose co-transporter 2 (SGLT2) inhibitors

NAME : DAPAGLIFLOZIN TABLETS 10 MG

AVAILABLE AS : PRIDAPA

PHARMACOLOGICAL ACTION : By inhibiting SGLT2, dapagliflozin blocks reabsorption of filtered glucose in the kidney, increasing urinary glucose excretion and reducing blood glucose levels. Its mechanism of action is independent of pancreatic β cell function and modulation of insulin sensitivity

USED FOR : PRIDAPA (Dapagliflozin) is mainly used to treat type 2 diabetes. It can also be used to treat heart failure and chronic kidney disease (CKD). PRIDAPA (Dapagliflozin) is usually prescribed if: you have type 2 diabetes and cannot take metformin.

SIDE EFFECTS : Cough.
difficulty with swallowing.
dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position.
dry mouth.
fainting.
increase in heart rate.
large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals.
lightheadedness.

WARNINGS : PRIDAPA (Dapagliflozin) can sometimes cause painful urination or pink, or bloody urine that are usually signs of a urinary tract infection, but they may also be signs of a rare condition (bladder cancer). Bladder cancer may or may not be caused by PRIDAPA (Dapagliflozin) Tell your doctor right away if you develop these symptoms.

OUR ADVICE : DO NOT SELF MEDICATE

This is a computer generated document.   Print Date : 30/09/2025 06:56 AM (IST)